THE CANADIAN ENTOMOLOGIST

Index to Volume 132 (2000)

THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

Agosti D. See Grimaldi and Agosti, 691
Ahn K-J. See Maruyama and Ahn, 567
Aldrich JR, Zhang A, Oliver JE. Attractant pheromone and allomone from the metathoracic scent gland of a broad-headed bug (Hemiptera: Alydidae), 915
Allen DC. See Silk *et al.*, 681
Anderson P. See Schlyter *et al.*, 965
Apigian K, Wheelwright NT. Forest ground beetles (Coleoptera: Carabidae) on a boreal island: habitat preferences and the effect of experimental removals, 627
Aragón A, Morón M-A. Description of third-instar larva of two species of *Phyllophaga* (Coleoptera: Melolonthidae), 323
Barker PS. See Lamb *et al.*, 591
Barrett JW, Ladd TR, Primavera MJ, Retnakaran A, Sohi SS, Palli SR. Nucleopolyhedrovirus pathology in spruce budworm larvae, 581
Bates SL, Borden JH, Savoie A, Blatt SE, Lait CG, Kermode AR, Bennett RG. Impact of feeding by *Leptoglossus occidentalis* (Hemiptera: Coreidae) on the major storage reserves of mature Douglas-fir (Pinaceae) seeds, 91
Beaver RA. Ambrosia beetles (Coleoptera: Platypodidae) of the South Pacific, 755
Behan-Pelletier VM. Ceratozetidae (Acari: Oribatida) of arboreal habitats, 153
Behringer A. See Vernon *et al.*, 231
Bennett R. See Miller *et al.*, 243
Bennett RG. See Bates *et al.*, 91
Berisford CW. See Birgersson *et al.*, 951
See Orr *et al.*, 373
Birgersson G. See Schlyter *et al.*, 965
Birgersson G, Dalusky MJ, Berisford CW. Identification of an aggregation pheromone for *Pityogenes hopkinsi* (Coleoptera: Scolytidae), 951
Bittner LA. See Pree *et al.*, 69
Blatt SE. See Bates *et al.*, 91
Bohlmann J. See Seybold *et al.*, 697
Boisvert JL. See Tessier *et al.*, 59
Borden JH. See Bates *et al.*, 91
See Huber *et al.*, 649
See Miller and Borden, 183
See Miller *et al.*, 243
See Poirier and Borden, 81
Bosco D. See Demichelis *et al.*, 519
Bouchard P, Wheeler TA, Goulet H. Design for a low-cost, covered, ramp pitfall trap, 387
Brooks JE. See Maclauchlan and Brooks, 773
Brown A. See Poinar *et al.*, 695
Browne LE. See Stephen and Browne, 983
Brunelle P-M. A new species of *Neurocordulia* (Odonata: Anisoptera: Corduliidae) from eastern North America, 39
Byers JA. See Schlyter *et al.*, 965
Caciagli P. See Demichelis *et al.*, 519
Chen X. See McElfresh *et al.*, 775
Cook J. Four new species of *Bdelyrus* (Coleoptera: Scarabaeidae: Scarabaeinae) and a redescription of *Bdelyrus lagopus*, 551
Cook J, Peck SB. Aphodiinae (Coleoptera: Scarabaeidae) of the Galápagos Islands, 281
Cossentine JE, Kuhlmann U. Status of *Ageniaspis fuscicollis* (Hymenoptera: Encrytidae), an introduced parasitoid of the apple ermine moth (Lepidoptera: Yponomeutidae), 685
Cox-Foster D. See Garcia-Salazar *et al.*, 1
Cusson M, Palli SR. Can juvenile hormone research help rejuvenate integrated pest management?, 263
Dallara PL, Seybold SJ, Meyer H, Tolasch T, Francke W, Wood DL. Semiochemicals from three species of *Pityophthorus* (Coleoptera: Scolytidae): identification and field response, 889
Dalusky MJ. See Birgersson *et al.*, 951
DeBarr GL. See de Groot and DeBarr, 843
See Orr *et al.*, 373
See Rappaport *et al.*, 925
DeBarr GL, Hanula JL, Niwa CG, Nord JC. Synthetic pheromones disrupt male *Dioryctria* spp. moths in a loblolly pine seed orchard, 345
de Groot P. See Miller *et al.*, 243
See Rappaport *et al.*, 925

de Groot P, DeBarr GL. Response of cone and twig beetles (Coleoptera: Scolytidae) and a predator (Coleoptera: Cleridae) to pityol, conophthorin, and verbenone, 843
Dedes J. See Massé *et al.*, 573
See van Frankenhuyzen and Dedes, 341
See van Frankenhuyzen *et al.*, 505
del Rio Mora AA. See Rappaport *et al.*, 925
Demichelis S, Bosco D, Manino A, Marian D, Caciagli P. Distribution of *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotypes in Italy, 519
Fields PG, Korunic Z. Diatomaceous earth to control *Cryptolestes ferrugineus* (Coleoptera: Cucujidae) in stored barley in farm granaries, 247
Fitzpatrick SM, Troubridge JT, Henderson D. *Ochropleura implecta* (Lepidoptera: Noctuidae), a new cutworm pest of cranberries, 365
Fleischer SJ. See Garcia-Salazar *et al.*, 1
Francke W. See Dallara *et al.*, 889
Galloway TD. See Pilgrim and Galloway, 15
Garcia-Salazar C, Gildow FE, Fleischer SJ, Cox-Foster D, Lukezic FL. Alimentary canal of adult *Acalymma vittata* (Coleoptera: Chrysomelidae): morphology and potential role in survival of *Erwinia tracheiphila* (Enterobacteriaceae), 1
Gibson GAP. Differentiation of the species of *Urolepis* (Hymenoptera: Chalcidoidea: Pteromalidae), potential biocontrol agents of filth flies (Diptera: Muscidae), 391
Gilbert N, Raworth DA. Insects and temperature—differential effects of experimental conditions on growth and development, 539
Gildow FE. See Garcia-Salazar *et al.*, 1
Goulet H. See Bouchard *et al.*, 387
Gries G. See Mistal *et al.*, 867
Gries R. See Huber *et al.*, 649
Grimaldi D, Agosti D. The oldest ants are Cretaceous, not Eocene: comment, 691
Guillet S, Josselin N, Vancassel M. Multiple introductions of the *Forficula auricularia* species complex (Dermaptera: Forficulidae) in eastern North America, 49
Haack RA. See Poland *et al.*, 853
Hamilton KGA. Five genera of new-world "shovel-headed" and "spoon-bill" leafhoppers (Hemiptera: Cicadellidae: Dorycephalini and Hecalini), 429
Hanks LM. See Paine *et al.*, 907
Hanula JL. See DeBarr *et al.*, 345
Harada T, Spence JR. Nymphal density and life histories of two water striders (Hemiptera: Gerridae), 353
Henderson D. See Fitzpatrick *et al.*, 365
Hoover SER. See Lindgren *et al.*, 877
Huber DPW, Borden JH, Jeans-Williams NL, Gries R. Differential bioactivity of conophthorin on four species of North American bark beetles (Coleoptera: Scolytidae), 649
Hunt DWA, Tan CS. Overwintering densities and survival of the Colorado potato beetle (Coleoptera: Chrysomelidae) in and around tomato (Solanaceae) fields, 103
Hunter FF. See Ossowski and Hunter, 213
Jeans-Williams NL. See Huber *et al.*, 649
See Miller *et al.*, 243
Johnson E. See Negrón *et al.*, 119
Johnson NF. A new species and revision of the concept of the genus *Thoronidea* (Hymenoptera, Platygastroidea: Scelionidae), 301
Josselin N. See Guillet *et al.*, 49
Kabaluk T. See Vernon *et al.*, 231
Keeling CI. See Lindgren *et al.*, 877
Kermode AR. See Bates *et al.*, 91
Korunic Z. See Fields and Korunic, 247
Kudo S-I, Mori S. Ploidy does not affect fluctuating asymmetry in the carpenter bee (Hymenoptera: Anthophoridae), 333
Kuhlmann U. See Cossentine and Kuhlmann, 685
Lacoursière JO. See Tessier *et al.*, 59
Ladd TR. See Barrett *et al.*, 581
Lait CG. See Bates *et al.*, 91
Lamb, RJ. See Wise and Lamb, 369
Lamb RJ, McKenzie RIH, Wise IL, Barker PS, Smith MAH, Olfert OO. Resistance to *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) in spring wheat (Gramineae), 591
Lamb RJ, Tucker JR, Wise IL, Smith MAH. Trophic interaction between *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) and spring wheat: implications for yield and seed quality, 607

Lanier GN. See Phillips and Lanier, 811
Lindgren BS, Hoover SER, MacIssac AM, Keeling CI, Slessor KN. Lineatin enantiomer preference, flight periods, and effect of pheromone concentration and trap length on three sympatric species of *Trypodendron* (Coleoptera: Scolytidae), 877
Linton DA. See Safranyik *et al.*, 799
Lobo JM. Species diversity and composition of dung beetle (Coleoptera: Scarabaeoidea) assemblages in North America, 307
Löfqvist J. See Schlyter *et al.*, 965
Lonergan GC. See Silk *et al.*, 681
Lukezic FL. See Garcia-Salazar *et al.*, 1
Macías-Sámano JE. See Moser and Macías-Sámano, 765
MacIssac AM. See Lindgren *et al.*, 877
Mackauer M. See Völkl and Mackauer, 197
Maclauchlan LE, Brooks JE. Temperature requirements for *Pissodes terminalis* (Coleoptera: Curculionidae), 773
Manino A. See Demichelis *et al.*, 519
Marian D. See Demichelis *et al.*, 519
Maruyama M, Ahn K-J. Redescription of *Liparocephalus litoralis*, and key to the species of *Liparocephalus* (Coleoptera: Staphylinidae: Aleocharinae), 567
Massé A, van Frankenhuyzen K, Dedes J. Susceptibility and vulnerability of third-instar larvae of the spruce budworm (Lepidoptera: Tortricidae) to *Bacillus thuringiensis* subsp. *kurstaki*, 573
McCravy KW. See Orr *et al.*, 373
McElfresh JS, Chen X, Ross DW, Millar JG. Sex pheromone blend of the pandora moth (Lepidoptera: Saturniidae), an outbreak pest in pine forests (Pinaceae), 775
McKenzie RIH. See Lamb *et al.*, 591
Meyer H. See Dallara *et al.*, 889
Millar JG. See McElfresh *et al.*, 775
See Paine *et al.*, 907
Miller DR. Vertical displacement of *Ips latidens* and *Ips pini* (Coleoptera: Scolytidae) by semiochemical interruption, 789
Miller DR, Borden JH. Dose-dependent and species-specific responses of pine bark beetles (Coleoptera: Scolytidae) to monoterpenes in association with pheromones, 183
Miller DR, Pierce HD Jr, de Groot P, Jeans-Williams N, Bennett R, Borden JH. Sex pheromone of *Conophthorus ponderosae* (Coleoptera: Scolytidae) in a coastal stand of western white pine (Pinaceae), 243
Mistal C, Takács S, Gries G. Evidence for sonic communication in the German cockroach (Dictyoptera: Blattellidae), 867
Monobrullah MD, Nagata M. Effects of larval age on susceptibility of *Spodoptera litura* (Lepidoptera: Noctuidae) to *Spodoptera litura* multiple nuclear polyhedrosis virus, 337
Mori S. See Kudo and Mori, 333
See Rappaport *et al.*, 925
Morón M-A. See Aragón and Morón, 323
Moser JC, Macías-Sámano JE. Tarsonemid mite associates of *Dendroctonus frontalis* (Coleoptera: Scolytidae): implications for the historical biogeography of *D. frontalis*, 765
Muza A. See Nagarkatti *et al.*, 259
Nagarkatti S, Muza A, Saunders M. Meridic diet for *Endopiza viteana* (Lepidoptera: Tortricidae), 259
Nagata M. See Monobrullah and Nagata, 337
Negrón JF, Schaupp WC, Johnson E. Development and validation of a fixed-precision sequential sampling plan for estimating brood adult density of *Dendroctonus pseudotsugae* (Coleoptera: Scolytidae), 119
Niwa CG. DeBarr *et al.*, 345
Nord JC. DeBarr *et al.*, 345
Nystrom C. See van Frankenhuyzen *et al.*, 505
Olfert OO. See Lamb *et al.*, 591
Oliver JE. See Aldrich *et al.*, 915
Onstad DW. See Poland *et al.*, 853
Orr DB, Suh CP-C, McCravy KW, Berisford CW, DeBarr GL. Evaluation of inundative releases of *Trichogramma exiguum* (Hymenoptera: Trichogrammatidae) for suppression of Nantucket pine tip moth (Lepidoptera: Tortricidae) in pine (Pinaceae) plantations, 373
Ossowski A, Hunter FF. Distribution patterns, body size, and sugar-feeding habits of two species of *Chrysops* (Diptera: Tabanidae), 213
Ostaff DP, Quiring DT. Population trends of a specialist herbivore, the spruce bud moth, in young white spruce stands, 825
Paine EO. See Paine *et al.*, 907

Paine TD, Hanks LM, Millar JG, Paine EO. Attractiveness and suitability of host tree species for colonization and survival of *Phoracantha semipunctata* (Coleoptera: Cerambycidae), 907
Palli SR. See Barrett *et al.*, 581
See Cusson and Palli, 263
Peck SB. See Cook and Peck, 281
Pérusse JR, Wheeler TA. Revision of the nearctic species of *Lauxania* (Diptera: Lauxaniidae), 411
Petrice TR. See Poland *et al.*, 853
Phillips TW, Lanier GN. Host specificity in *Pissodes strobi* (Coleoptera: Curculionidae): roles of geography, genetics, and behavior, 811
Pierce HD Jr. See Miller *et al.*, 243
Pilgrim RLC, Galloway TD. Descriptions of flea larvae (Siphonaptera: Ceratophyllidae: *Ceratophyllus* spp.) found in the nests of swallows (Aves: Passeriformes: Hirundinidae) in North America, north of Mexico, 15
Plowright CMS, Silverman A. Nectar and pollen foraging by bumble bees (Hymenoptera: Apidae): choice and tradeoffs, 677
Poinar G, Urbani CB, Brown A. The oldest ants are Cretaceous, not Eocene: reply, 695
Poirier LM, Borden JH. Influence of diet on repellent and feeding-deterrent activity of larval oral exudate in spruce budworms (Lepidoptera: Tortricidae), 81
Poland TM, Haack RA, Petrice TR, Sadof CS, Onstad DW. Dispersal of *Tomicus piniperda* (Coleoptera: Scolytidae) from operational and simulated mill yards, 853
Powell GW, Wikeem BM, Sturko A. Biology of *Agapeta zoegana* (Lepidoptera: Cochylidae), propagated for the biological control of knapweeds (Asteraceae), 223
Pree DJ, Bittner LA, Whitty KJ. Field bioassay methodologies for the detection of resistance to acaricides in populations of European red mite (Acari: Tetranychidae), 69
Primavera MJ. See Barrett *et al.*, 581
Quednau FW. Taxonomic notes on the genus *Monelliopsis* (Hemiptera: Aphididae: Calaphidinae) with descriptions of three new species, 135
Quiring DT. See Ostaff and Quiring, 825
Raffa KF. See Seybold *et al.*, 697
Rappaport NG, Stein JD, del Rio Mora AA, DeBarr G, de Groot P, Mori S. Responses of *Conophthorus* spp. (Coleoptera: Scolytidae) to behavioral chemicals in field trials: a transcontinental perspective, 925
Raworth DA. See Gilbert and Raworth, 539
Retnakaran A. See Barrett *et al.*, 581
Robertson IC. Reproduction and developmental phenology of *Ips perturbatus* (Coleoptera: Scolytidae) inhabiting white spruce (Pinaceae), 529
Ross DW. See McElfresh *et al.*, 775
Ryall KL, Smith SM. Brood production and shoot feeding by *Tomicus piniperda* (Coleoptera: Scolytidae), 939
Sadof CS. See Poland *et al.*, 853
Safranyik L, Linton DA, Shore TL. Temporal and vertical distribution of bark beetles (Coleoptera: Scolytidae) captured in barrier traps at baited and unbaited lodgepole pines the year following attack by the mountain pine beetle, 799
Saunders M. See Nagarkatti *et al.*, 259
Savoie A. See Bates *et al.*, 91
Schaupp WC. See Negrón *et al.*, 119
Schlyter F, Zhang Q-H, Anderson P, Byers JA, Wadhams LJ, Löfqvist J, Birgersson G. Electrophysiological and behavioural responses of *Tomicus piniperda* and *Tomicus minor* (Coleoptera: Scolytidae) to non-host leaf and bark volatiles, 965
Schmaedick MA, Shelton AM. Arthropod predators in cabbage (Cruciferae) and their potential as naturally occurring biological control agents for *Pieris rapae* (Lepidoptera: Pieridae), 655
Seybold SJ. See Dallara *et al.*, 889
Seybold SJ, Bohlmann J, Raffa KF. Biosynthesis of coniferophagous bark beetle pheromones and conifer isoprenoids: evolutionary perspective and synthesis, 697
Shelton AM. See Schmaedick and Shelton, 655
Shore TL. See Safranyik *et al.*, 799
Shorthouse JD. See St John and Shorthouse, 635
Silk PJ, Lonergan GC, Allen DC, Spear-O'Mara J. Potential sex pheromone components of the saddled prominent (Lepidoptera: Notodontidae), 681
Silverman A. See Plowright and Silverman, 677
Slessor KN. See Lindgren *et al.*, 877
Smith MAH. See Lamb *et al.*, 591
See Lamb *et al.*, 607
Smith SM. See Ryall and Smith, 939

Sohi SS. See Barrett *et al.*, 581
Spear-O'Mara J. See Silk *et al.*, 681
Spence JR. See Harada and Spence, 353
St John MG, Shorthouse JD. Allocation patterns of organic nitrogen and mineral nutrients within stem galls of *Diplolepis spinosa* and *Diplolepis triforma* (Hymenoptera: Cynipidae) on wild roses (Rosaceae), 635
Stein JD. See Rappaport *et al.*, 925
Stephen FM, Browne LE. Application of Eliminade™ parasitoid food to boles and crowns of pines (Pinaceae) infested with *Dendroctonus frontalis* (Coleoptera: Scolytidae), 983
Sturko A. See Powell *et al.*, 223
Suh CP-C. See Orr *et al.*, 373
Takács S. See Mistal *et al.*, 867
Tan CS. See Hunt and Tan, 103
Tessier L, Boisvert JL, Vought LB-M, Lacoursière JO. Characterization of *Hydropsyche slossonae* (Trichoptera: Hydropsychidae) capture net polypeptides, 59
Tolasch T. See Dallara *et al.*, 889
Trimble RM, Tyndall CA. Disruption of mating in the spotted tentiform leafminer (Lepidoptera: Gracillariidae) using synthetic sex pheromone, 107
Troubridge JT. See Fitzpatrick *et al.*, 365
Tucker JR. See Lamb *et al.*, 607
Tyndall CA. See Trimble and Tyndall, 107
Urbani CB. See Poinar *et al.*, 695
Vancassel M. See Guillet *et al.*, 49
van Frankenhuyzen K. See Massé *et al.*, 573
van Frankenhuyzen K, Dedes J. Exposing larvae of the spruce budworm (Lepidoptera: Tortricidae) to sublethal doses of *Bacillus thuringiensis* subsp. *kurstaki* does not affect survival during the pupal stage, 341
van Frankenhuyzen K, Nystrom C, Dedes J. Mortality, feeding inhibition, and recovery of spruce budworm (Lepidoptera: Tortricidae) larvae following aerial application of a high-potency formulation of *Bacillus thuringiensis* subsp. *kurstaki*, 505
Vernon RS, Kabaluk T, Behringer A. Movement of *Agriotes obscurus* (Coleoptera: Elateridae) in strawberry (Rosaceae) plantings with wheat (Gramineae) as a trap crop, 231
Völkl W, Mackauer M. Oviposition behaviour of aphidiine wasps (Hymenoptera: Braconidae, Aphidiinae): morphological adaptations and evolutionary trends, 197
Vought LB-M. See Tessier *et al.*, 59
Wadhams LJ. See Schlyter *et al.*, 965
Wheeler TA. See Bouchard *et al.*, 387
See Pérusse and Wheeler, 411
Wheelwright NT. See Apigian and Wheelwright, 627
Whitty KJ. See Pree *et al.*, 69
Wikeem BM. See Powell *et al.*, 223
Wise IL. See Lamb *et al.*, 591
See Lamb *et al.*, 607
Wise IL, Lamb RJ. Seasonal occurrence of plant bugs (Hemiptera: Miridae) on oilseed flax (Linaceae) and their effect on yield, 369
Wood DL. See Dallara *et al.*, 889
Zhang A. See Aldrich *et al.*, 915
Zhang Q-H. See Schlyter *et al.*, 965